

Open Research Online

The Open University's repository of research publications and other research outputs

Quantitative genetics of body size evolution on islands: an individual-based simulation approach

Journal Item

How to cite:

Diniz-Filho, JAF; Jardim, L; Rangel, TF; Holden, Philip; Edwards, Neil; Hortal, J; Santos, AMC and Raia, P (2019). Quantitative genetics of body size evolution on islands: an individual-based simulation approach. *Biology Letters*, 15(10)

For guidance on citations see [FAQs](#).

© 2019 The Authors



<https://creativecommons.org/licenses/by-nc-nd/4.0/>

Version: Accepted Manuscript

Link(s) to article on publisher's website:
<http://dx.doi.org/doi:10.1098/rsbl.2019.0481>

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

oro.open.ac.uk

Quantitative Genetics of Body Size Evolution in Islands: an individual-based simulation approach

Journal:	<i>Biology Letters</i>
Manuscript ID	RSBL-2019-0481.R1
Article Type:	Research
Date Submitted by the Author:	n/a
Complete List of Authors:	Diniz-Filho, José Alexandre; Universidade Federal de Goiás, Ecologia Jardim, Lucas; Universidade Federal de Goiás, Ecologia Rangel, Thiago; Universidade Federal de Goiás, Departamento de Ecologia, ICB Holden, Philip; The Open University, Environment, Earth and Ecosystem Sciences Edwards, Neil; The Open University, Environment Earth and Ecosystems Hortal, Joaquín; Museo Nacional de Ciencias Naturales, Biodiversity and Global Change Lab Santos, Ana; Centro de Biologia Ambiental , Faculdade de Ciências da Universidade de Lisboa Raia, Pasquale; Università degli Studi di Napoli Federico II, Dipartimento di Scienze della Terra, dell'Ambiente e delle Risorse
Subject:	Ecology < BIOLOGY, Evolution < BIOLOGY, Palaeontology < BIOLOGY
Categories:	Evolutionary Biology
Keywords:	adaptation, body size, island rule, natural selection, Homo floresiensis, dwarfing

Author-supplied statements

Relevant information will appear here if provided.

Ethics

Does your article include research that required ethical approval or permits?:

This article does not present research with ethical considerations

Statement (if applicable):

CUST_IF_YES_ETHICS :No data available.

Data

It is a condition of publication that data, code and materials supporting your paper are made publicly available. Does your paper present new data?:

My paper has no data

Statement (if applicable):

CUST_IF_YES_DATA :No data available.

Conflict of interest

I/We declare we have no competing interests

Statement (if applicable):

CUST_STATE_CONFLICT :No data available.

Authors' contributions

This paper has multiple authors and our individual contributions were as below

Statement (if applicable):

JAFD-F, P. R., J. H. and A. M. S. conceived the study and wrote the first draft. JAFD-F, L.J. and TFR wrote the main codes and L.J. conducted all simulations; JAFD-F, J. H., A.M.S. and PR discussed the results and interpretations, and P.H.H. and N.R.E. provided original environmental data for time series analyses. All authors contributed to writing of the final version of the manuscript.

To: *Biology Letters* (Research Article)

Section: *Evolutionary Biology*

Quantitative Genetics of Body Size Evolution in Islands: an individual-based simulation approach

José Alexandre F. Diniz-Filho^{1*}, Lucas Jardim², Thiago F. Rangel¹, Phillip B. Holden³, Neil R. Edwards³, Joaquín Hortal^{1,4,6}, Ana M. C. Santos^{5,6}, Pasquale Raia⁷

1. *Departamento de Ecologia, ICB, Universidade Federal de Goiás (UFG)*

2. *INCT EECBio, DTI program*

3. *Environment, Earth and Ecosystem Sciences, The Open University, UK*

4. *Department of Biogeography and Global Change, Museo Nacional de Ciencias Naturales (MNCN-CSIC), C/José Gutiérrez Abascal 2, 28006 Madrid, Spain*

5. *GLOCEE - Global Change Ecology and Evolution Group, Departamento de Ciencias de la Vida, Universidad de Alcalá, 28805 Alcalá de Henares, Spain*

6. *cE3c – Centre for Ecology, Evolution and Environmental Changes, Faculdade de Ciências, Universidade de Lisboa. Campo Grande, Edifício C2, Piso 5, 1749-016 Lisboa, Portugal*

7. *Department DiSTAR, University of Naples Federico II, Via Cintia 21, 20126, Napoli, Italy*

* Corresponding author: diniz@ufg.br

ABSTRACT

According to the Island Rule small-bodied vertebrates will tend to evolve larger body size on island, whereas the opposite happens to large-bodied species. This controversial pattern has been studied at the macroecological and biogeographical scales, but new developments in quantitative evolutionary genetics now allow studying the Island Rule from a mechanistic perspective. Here we develop a simulation approach based on an Individual-Based Model (IBM) to model body size change on islands as a progressive adaptation to a moving optimum, determined by density-dependent population dynamics. We applied the model to evaluate body size differentiation in the pigmy extinct hominin *Homo floresiensis*, showing that dwarfing may have occurred in only about 360 generations (95% CI ranging from 150 to 675 generations). This result agrees with reports suggesting rapid dwarfing on large mammals on islands, as well as with the recent discovery that small-sized hominins lived in Flores as early as 700 kyr ago. Our simulations illustrate the power of analyzing ecological and evolutionary patterns from an explicit quantitative genetics perspective.

Keywords: adaptation, body size, dwarfing, island rule, *Homo*

1. Introduction

The Island rule is amongst the most widely discussed body size patterns in Ecology, with a long history of heated debates and controversies [1-5]. This biogeographical pattern states that after colonizing an island, small-bodied vertebrates will tend to evolve towards larger bodies (i.e. gigantism), whereas large-bodied species will reduce their body size (i.e. dwarfing) [6,7], in a continuous linear grade [8]. Despite the enduring controversy and the discussions around the ubiquity of the pattern and the ecological and evolutionary mechanisms behind it [2, 5, 9], islands are legitimately renowned for some spectacular examples of dwarfing in large artiodactyls, proboscideans and even hominids [see 10-14].

The Island rule was traditionally studied at the macroecological and biogeographical scales, comparing several island species at once and trying to correlate their shift in body size from their presumed ancestor [8] with i) the islands physical and environmental characteristics or ii) with biotic characteristics that could be surrogates of competition or predation [15-18]. At the macroevolutionary scale these patterns are better described by a punctuated model in which body size shifts occur fast after speciation and are followed by a stasis [19]. However, regardless of the macroevolutionary and macroecological generalization of the Island rule, it is important to recognize that this pattern must derive from population-level processes that are still little explored.

Body size changes in insular populations may occur over a relatively short time, involving combined effects of genetic adaptations and phenotypic plasticity [16, 20-21]. Palovacs [20] proposed a general life-history framework to explain the evolutionary trajectory of body size on islands (see also [22] for first insights on life-history patterns in island species evolution). Both gigantism and dwarfing would thus be explained by a complex interaction of adaptive responses involving the balance between reduction of growth rates under resource depletion and reduced mortality on islands [23], or simply by changes in life history traits related to reduction in the age of sexual maturity [24, 25]. Ecological processes involving phenotypic plasticity and life-history shifts may occur fast, in the first few generations after the island colonization event, triggering or canalizing long-term changes by natural selection due to intraspecific competition [16, 24].

Several theoretical models have been proposed to describe the evolutionary dynamics of quantitative traits under different processes since first developments of quantitative and population genetics in early 20th century [26]. In the context of insular evolution, Diniz-Filho and Raia [27] used quantitative evolutionary genetics models to evaluate brain and body size differentiation in *Homo floresiensis*, showing that adaptive explanations for its dwarfing from an *H. erectus* ancestor are plausible. However, incorporating complex evolutionary dynamics with more realistic population dynamics under demographic and environmental stochasticity requires alternative approaches. Here we expand upon our previous analysis [27] by using a simulation approach based on an individual-based model (IBM) grounded on first principles of population and quantitative genetics [26, 28]. Our main focus is to evaluate the plausibility of fast dwarfism in *H. floresiensis* [29] incorporating several evolutionary processes and based on a new model in which adaptation occurs under a moving optimum determined by density-dependent population dynamics.

2. Methods

(a) *Modeling evolutionary processes within and between generations*

The details of the simulation model and the parametrization are available as Electronic Supplementary Material (ESM). The simulation starts with a vector **G** of genotypic values for body size from N_0 individuals representing the initial deme colonizing the island, sampled from a normal distribution with mean u_G and additive genetic variance v_A , for a single quantitative trait (i.e., body size), thus assuming an infinitesimal model [30]. After island colonization, the deme starts to grow in numbers following a logistic process up to the island carrying capacity K . The process stops after t_G generations, when the mean phenotypic value is close to the final adaptive peak defined for the island.

Phenotypic values of the population (**P**) are obtained by adding to **G** a vector with randomly distributed environmental effects **E** with mean zero and environmental variance v_E (which is in turn determined assuming a known heritability h^2 for the trait), with phenotypic plasticity incorporated as a linear reaction norm [31-34]. Fitness W_i for each individual is defined according to a Gaussian approximation of stabilizing selection with an

101 optimum equal to O (the adaptive peak; see below for properties of this optimum). In our
102 model, natural selection acts through both differential survival (i.e., before reproduction)
103 and fecundity (i.e., number of offspring). Inbreeding is modeled by a Markovian
104 approximation tracking brother-sister breeding and reduces fitness by increasing mortality
105 in newborns [35, 36].

106 Variance in populations is restored by incorporating mutation and migration to our
107 model. Mutation is added as a mutation kernel [37], adding to \mathbf{G} values of the newborn a
108 vector with mean 0 and variance v_U/v_A (assumed to vary between $0.02-0.03v_A$ for polygenic
109 traits such as body size [38]). We also added to the model the probability of island
110 colonization by new individuals, with genotypic values sampled from the original values of
111 \mathbf{G} at the beginning of the simulations (assuming that the continental population is large and
112 under demographic and genetic equilibrium).

113 Lastly, under a classical adaptive model for islands the optimum body size O would
114 be smaller than on the continent (so that the Gaussian adaptive landscape would converge
115 to a Gaussian, non-linear directional selection function if mean \mathbf{P} differs from O [39, 40]).
116 It is realistic that the intensity of dwarfing should be density-dependent and would thus be
117 small in the initial phases of colonization (i.e. when $N \ll K$). For instance, a reduction in
118 the age of sexual maturity due to abundant per-capita resources leading to small body size
119 would be initially attributed to plasticity [16, 23, 31] and followed by a second phase of
120 intraspecific competition and selection (this pattern also appears in Lister's [16] two-phase
121 model for dwarfing). Thus, it is expected that the initial adaptive peak (O_0) equals the mean
122 continental body size at the beginning, and natural selection would start favoring smaller
123 individuals as the population starts growing towards K .

124 At each step of the simulation individual body size will thus evolve and the time to
125 adaptation is defined as the time taken by mean \mathbf{P} to achieve the final optimum. As this is
126 an iterative process across generations, it is possible to record several parameters and
127 outcomes of the model at intermediate steps, including the mean and variance of \mathbf{G} ,
128 population density, inbreeding level, realized heritability h^2 , as well as statistics describing
129 the intensity of selection process (i.e., mean-standardized selection gradient, [41, 42]).

(b) *Evolution of body size in Homo floresiensis*

We parametrized and tested our simulation model using empirical data for *Homo floresiensis*, a case previously investigated by two of us [27]. *H. floresiensis* was a small-bodied fossil hominin of about 25-30 kg discovered on Flores Island, Indonesia, in 2004 [11, 43]. It most likely represents a dwarf form of *Homo erectus*, the first hominine species to colonize South-East Asia almost 2 mya [43, 44].

We simulated dwarfing in *H. floresiensis* from a large-bodied ancestor in the range of *H. erectus* (i.e., 45-55 kg), under 10,000 random combinations of parameters sampled within the ranges defined in Table 1 (see also ESM and Diniz-Filho and Raia [27] for details and references). Adaptation to the new final peak is considered “successful” if mean \bar{G} is below 30 kg, given the uncertainty of around 5% around the 27 kg usually estimated for *H. floresiensis* [11, 43-44]. Body mass values are used here as a general surrogate for body size, in a comparative and interspecific sense [e.g., 45]. The main response variable in our simulations is the time for adaptation, t_G . Finally, we analyzed how the simulation parameters explain the realized mean-standardized selection gradient across simulations (see also Fig. S2 in ESM).

3. Results

Under the realistic range of demographic and genetic parameters successful adaptations happened in 97.3% of the simulations. In these simulations, the time to adaptation t_G is log-normally distributed, with median time equal to 346 generations, and 95% non-parametric confidence intervals ranging from 150 to 675 generations (Fig. 1). This median corresponds to about 5190 years (95% CI ranging from 2250 and 10125 years), assuming some 15 years as the mean age for sexual maturity and generation length (see also ESM for some additional results fixing some of the most important parameters in the simulations).

We explained about 70% of the variation in log-transformed t_G across the simulations by a multiple linear regression. Lower t_G are found for simulations with high mutation rate, large initial population size and large size of the demes recolonizing the

island (even with maximum of 10 new individuals with a maximum low probability of recolonizing equal to 0.1 per generation). Despite the high overall explanatory power of this model and the high effect size of these three parameters, the individual ability of each parameter to predict t_G is rather low (see ESM Fig. S4), suggesting that the response emerge from a complex interaction of different parameters (which may actually explain log-normally distribution of t_G)

Realized mean-standardized selection gradients in the simulation are left-skewed (Fig. 2), with median equal to -0.29 (95% CI ranging from -0.167 to -0.562), with about 82% of its variance explained by simulation parameters. Higher selection strength is necessary to drive adaptation in either shorter adaptive landscapes, when ancestral mean body sizes are large, and/or when the amount of variation explained by phenotypic plasticity is low (although these two last effects are smaller; see ESM Table 2).

4. Discussion

Our simulations support the hypothesis that *H. floresiensis* plausibly became dwarf over a relatively short time. Gomez-Robles [29] recently raised the question of whether it would be possible for the extreme reduction of the brain and body of *H. floresiensis* to have occurred over a mere 300,000 years, given estimated ages of *H. floresiensis* fossil record [43, 46-47]. Confirming the results from classical quantitative genetics models [27], we showed here that dwarfism in *H. floresiensis* could have been much faster than this. In 97% of the simulations, adaptation occurred with median time for dwarfism equal to some 5000 years, or ~350 generations. Indeed, it is documented that the effects of natural selection on trait variations can be very fast on islands, covering a few tenths of generations at most [19, 48-50]. Moreover, recent analyses point out to larger effects of phenotypic plasticity, especially in early phases of island colonization, as also proposed in Lister's [16] two-phase model and by acceleration of life-history traits evolution due to biotic release [20, 24-25].

As expected, the mean-standardized selection gradient is higher than previously found by using analytical solutions in ref. [27], at around -0.3, but even so this value is

quite close to common estimates in natural populations [51]. Moreover, these values emerged from a realistic set of demographic and genetic parameters and widely accepted processes. Although these coefficients are indisputably related to the adaptive landscape adopted, we assumed relatively weak selection within generations (i.e., $w^2 < 100v_A$), and a moving peak that allows population survival and quick expansion just after island colonization.

Of course, our analyses do not make it possible to establish that *H. floresiensis* is a dwarfed form of *H. erectus* [see 43, 52-54 for discussions], or to state that speciation occurred as fast as 5,000 years, as many other complex morphological traits, in addition to body size, differentiate *H. floresiensis* from *H. erectus* (and other related species; see [43, 52-54]). Rather, our analyses should be better viewed as a proof of concept that fast dwarfing is plausible even in hominins.

The simulation model used here still uses a phenomenological approach to quantitative genetics [55], in the sense that the adaptive landscape and peak are defined on the target dwarfed species evolving from a large-bodied ancestor. Even so, we believe our simulation illustrates the power of analyzing ecological and evolutionary patterns from an explicit quantitative genetics perspective. We argue the IBM model developed here could be viewed as a starting point for a quasi-experimental model, to better understand which demographic and genetic parameters may really lead to rapid dwarfing in insular vertebrates.

Acknowledgements

We thank three reviewers for their nice suggestions that improved original version of this manuscript. This paper is a contribution of the INCT in Ecology, Evolution and Biodiversity Conservation founded by MCTIC/CNPq/FAPEG (grant 465610/2014-5), arising from the workshop “Fast Evolution on Islands”, organized by AMCS and JAFD-F. JH was supported by the project ‘Predicting diversity variations across scales through process-based models linking community ecology and biogeography’ (CNPq PVE 314523/2014-6), and AMCS by an Spanish MICIU Juan de la Cierva-Incorporación grant.

REFERENCES

1. Lomolino MV. 2005. Body size evolution in insular vertebrates: generality of the island rule. *J. Biogeogr.* **32**, 1683-1699.
2. Meiri S, Raia P, Phillimore A. 2010. Slaying dragons: limited evidence for unusual body size evolution on islands. *J. Biogeogr.* **38**, 89-100.
3. Raia P, Carotenuto F, Meiri S. 2010. One size does not fit all: no evidence for an optimal body size on islands. *Glob. Ecol. Biogeogr.*, **19**, 475-484.
4. Faurby S, Svenning JC. 2016. Resurrection of the island rule: human-driven extinctions have obscured a basic evolutionary pattern. *Am. Nat.* **187**, 812-820.
5. Lokatis S, Jeschke JM. 2018. The island rule: An assessment of biases and research trends. *J. Biogeogr.* **45**, 289-303.
6. Foster, JB. 1964. Evolution of mammals on islands. *Nature* **202**, 234-235.
7. Van Valen, L. 1973. Pattern and balance of nature. *Evol. Theory* **1**, 31-49.
8. Lomolino MV. 1985. Body size of mammals on islands: the island rule reexamined. *Am. Nat.* **125**, 310-316.
9. Meiri S, Dayan T, Simberloff D. 2006. The generality of island rule reexamined. *J. Biogeogr.* **33**, 1571-1577.
10. van der Geer A, Lyras G, de Vos J, Dermitzakis M. 2012. *Evolution of island mammals: adaptation and extinction of placental mammals on islands*. Oxford, UK, Wiley-Blackwell.
11. Brown P, Sutikna MJ, Morwood MJ, Soejono RP, Jatmiko, Wayhu Saptomo E, Awe Due R.. 2004. A new small-bodied hominin from the Late Pleistocene of Flores, Indonesia. *Nature* **431**, 1055-1061.
12. Detroit F, Mijares AS, Corny J, Daver G, Zanolis C, Dizon E, Robles E, Grun R., Pipers PJ. 2019. A new species of Homo from the late pleistocene of the Phillipines. *Nature* **561**, 181-186.
13. Athanassiou A, Van der Geer AA, Lyras GA. 2019. Pleistocene insular Proboscidea of the Eastern Mediterranean: A review and update. *Quat. Sci. Rev.* **218**, 306-321.

- 247 14. Rozzi R. 2018. Space–time patterns of body size variation in island bovids: The key role of
248 predatory release. *J Biogeogr* **45**, 1196-1207.
- 249 15. Raia P, Meiri S. 2006. The island rule in large mammals: paleontology meets ecology. *Evolution*
250 **60**, 1731-1742.
- 251 16. Lister AM. 1996. Dwarfing in island elephants and deer: processes in relation to time of isolation.
252 In: Symposia of the Zoological Society of London (No. 69, pp. 277-292). London: The
253 Society, 1960-1999.
- 254
- 255 17. Lomolino MV, Sax DF, Palombo MR, van der Geer AA. 2012. Of mice and mammoths:
256 evaluations of causal explanations for body size evolution in insular mammals. *J. Biogeogr.*
257 **39**, 842-854.
- 258 18. McClain CR, Durst PA, Boyer AG, Francis CD. 2013. Unravelling the determinants of insular
259 body size shifts. *Biol. Lett.* **9**, 20120989.
- 260 19. Raia P, Meiri S. 2011. The tempo and mode of evolution: body sizes of island mammals. *Evolution*
261 **65**, 1927-1934.
- 262 20. Ozgul A, Tuljapurkar S, Benton TG, Pemberton JM, Clutton-Brock TH, Coulson T. 2009. The
263 dynamics of phenotypic change and the shrinking sheep of St. Kilda. *Science* **325**, 464-467.
- 264 21. Palovacs EP. 2003. Explaining adaptive shifts in body size on islands: a life history approach.
265 *Oikos* **103**, 37-44.
- 266 22. Case TJ. 1978. A general explanation for insular body size trends in terrestrial vertebrates.
267 *Ecology* **59**, 1-18.
- 268 23. Long ES, Courtney KS, Lippert JC, Wall-Scheffler CM. 2019. Reduced body size of insular
269 black-tailed deer is caused by slowed development. *Oecologia* **189**, 675-685.
- 270 24. Raia P, Barbera C, Conte M. 2003. The fast life of a dwarfed giant. *Evol. Ecol.* **17**, 293-312.
- 271 25. Migliano AB, Vinicius L, Lahr MM. 2012. Life history trade-offs explain the evolution of human
272 pygmies. *Proc. Natl. Acad. Sci.* **104**, 20216-20219.
- 273 26. Walsh B, Lynch M. 2018. *Evolution and selection of quantitative traits*. Oxford, UK: Oxford
274 University Press.
- 275 27. Diniz-Filho JAF, Raia P. 2017. Island Rule, quantitative genetics and brain–body size evolution
276 in *Homo floresiensis*. *Proc. R. Soc. Lond.* **284**, 20171065.

- 287 28. Roff D. 2010. *Modeling Evolution*. Oxford, UK: Oxford University Press.
- 288 29. Gómez-Robles A. 2016 The dawn of *Homo floresiensis*. *Nature* **534**, 188-189.
- 289 30. Barton N, Etheridge AM, Véber A. 2017. The infinitesimal model: Definition, derivation, and
290 implications. *Theor. Popul. Biol.* **118**, 50-73.
- 281 31. Sinclair ARE, Parkes JP. 2008. On being the right size: food-limited feedback on optimal body
282 size. *J. Anim. Ecol.* **77**, 635-637.
- 283 32. Lande R. 2009. Adaptation to an extraordinary environment by evolution of phenotypic plasticity
284 and genetic assimilation. *J. Evol. Biol.* **22**, 1435–1446.
- 285 33. Chevin, L. M., Lande, R., Mace, G. M. 2010. Adaptation, plasticity, and extinction in a changing
286 environment: towards a predictive theory. *PLoS Biol.* **8**, e1000357.
- 287 34. Schmid M, Guillaume F. 2017. The role of phenotypic plasticity on phenotypic differentiation.
288 *Heredity* **119**, 214–225.
- 289 35. Bittles AB, Neel JV. 1994. The cost of human inbreeding and their implications for variations at
290 the DNA level. *Nat. Genet.* **8**, 117-121.
- 291 36. Huisman J, Kruuk LEB, Ellis PA, Clutton-Brock T, Pemberton JM. 2016. Inbreeding depression
292 across the lifespan in a wild mammal population. *Proc. Natl. Acad. Sci.* **113**, 3585-3590.
- 293 37. Débarre F, Otto SP. 2016. Evolutionary dynamics of a quantitative trait in a finite asexual
294 population. *Theor. Popul. Biol.* **108**, 75–88.
- 295 38. Kemper KE, Visscher PM, Goddard ME. 2012. Genetic architecture of body size in mammals.
296 *Genome Biol.* **13**, 244.
- 297 39. Gomulkiewicz R, Holt RD. 1995. When does evolution by natural selection prevent extinction?
298 *Evolution* **49**, 201-204.
- 299 40. Chevin LM, Lande R. 2011. Adaptation to marginal habitats by evolution of increased phenotypic
300 plasticity. *J. Evol. Biol.* **24**, 1462-1476.
- 301 41. Kingsolver JG, Hoekstra HE, Hoekstra JM, Berrigan D, Vignieri SN, Hill CE, Hoang A, Gibert
302 P, Beerli P. 2001. The strength of phenotypic selection in natural populations. *Am. Nat.* **157**,
303 245-261.
- 304 42. Matsumura S, Arlinghaus R, Ieckman U. 2012. Standardizing selection strengths to study selection
305 in the wild: a critical comparison and suggestions for the future. *Bioscience* **62**, 1039-1054.

- 306 43. Baab KL. 2016b. The place of *Homo floresiensis* in human evolution. *J. Anthropol. Sci.* **94**, 5–
307 18.
- 308 44. Baab KL. 2016a. The role of neurocranial shape in defining the boundaries of an expanded *Homo*
309 *erectus* hypodigm. *J. Hum. Evol.* **92**, 1–21.
- 310 45. Grabowski M 2016. Bigger brains led to bigger bodies? The correlated evolution of human brain
311 and body size. *Curr. Anthropol.* **57**, 174–196.
- 312 46. van den Bergh GD, Kaifu Y, Kurniawan I, Kono RT, Brumm A, Setiyabudi E, Aziz F, Morwood
313 MJ. 2016. *Homo floresiensis* – like fossils from the early Middle Pleistocene of Flores. *Nature*
314 **534**, 245–248.
- 315 47. Dennell RW, Louys J, O'Regan HJ, Wilkinson DM. 2014. The origins and persistence of *Homo*
316 *floresiensis* on Flores: biogeographical and ecological perspectives. *Quat. Sci. Rev.* **96**, 98–
317 107.
- 318 48. Rozzi R Lomolino MV. 2017. Rapid Dwarfing of an Insular Mammal – The Feral Cattle of
319 Amsterdam Island. *Sci. Rep.* **7**, 8820.
- 320 49. Lister AM. 1989. Rapid dwarfing of red deer on Jersey in the last interglacial. *Nature* **342**, 539–
321 542.
- 322 50. Millien V. 2011. Mammals evolve faster in smaller islands. *Evolution* **65**, 1935–1944.
- 323 51. Hereford J, Hansen TF, Houle D. 2004. Comparing strengths of directional selection: How strong
324 is strong? *Evolution* **58**, 2133–2143.
- 325 52. Argue D, Groves CP, Lee MSY, Jungers WL. 2017. The affinities of *Homo floresiensis* based on
326 phylogenetic analyses of cranial, dental, and postcranial characters. *J. Hum. Evol.* **107**, 107–
327 133.
- 328 53. Dembo M, Matzke NJ, Mooers AO, Collard M. 2015. Bayesian analysis of a morphological
329 supermatrix sheds light on controversial fossil hominin relationships. *Proc. R. Soc. Lond.* **282**,
330 20150943.
- 331 54. Zeitoun V, Barriel V, Widiyanto H. 2016. Phylogenetic analysis of the calvaria of *Homo*
332 *floresiensis*. *CR PALEVOL* **15**, 555–568.
- 333 55. Stearns SC. 1983. *The Evolution of Life Histories*. Oxford, UK: Oxford University Press.

334
335

336

337

338 **Authors contributions** - JAFD-F, P. R., J. H. and A. M. S. conceived the study and wrote
339 the first draft. JAFD-F, L.J. and TFR wrote the main codes and L.J. conducted all
340 simulations; JAFD-F, J. H., A.M.S. and PR discussed the results and interpretations, and
341 P.H.H. and N.R.E. provided original environmental data for time series analyses. All
342 authors contributed to writing of the final version of the manuscript.

343 **Statement on Competing Interests** - We have no competing interests

344

345 **Ethics statement** - We declare all ethical procedures were followed during the current
346 study.

347 **Data accessibility statement** - All data necessary to reproduce the results in this paper can
348 be found in the electronic appendix

349

350

Figure Captions:

Fig. 1. Distribution of time for adaptation (in generations) obtained from the IBM simulating dwarfing of *H. floresiensis* based on 10,000 random combinations of parameters defined in Table 1. Median time is equal to 346 generations (CI 95% ranging from 150 and 675 generations).

Fig. 2. Distribution of mean-standardized selection gradients generated in the IBM simulating dwarfing of *H. floresiensis* based on 10,000 random combinations of parameters defined in Table 1. Median selection strength is equal to -0.29 (CI 95% ranging from -0.167 and -0.562).

For Review Only

Table 1. Genetic and demographic parameters used in the Individual-Based Model for body size evolution in *Homo floresiensis*.

<i>Parameter</i>	<i>Symbol</i>	<i>range of values</i>
Initial adaptive peak (kg)	O_0	45 – 55
Final adaptive peak (kg)	O	27
Heritability	h^2	0.6 – 0.85
Phenotypic coefficient of variation	Cv	0.04 – 0.06
mutational variance	v_U / v_A	0.02 – 0.04
Length of adaptive landscape (x v_A)	w^2	100 – 150
Phenotypic plasticity	b	0.1 – 0.5
Inbreeding depression	I	0.7
Initial inbreeding coefficient	F	0.1
Initial Population	N_0	25 – 100
Carrying capacity	K	10,000 x NPP_t
Number of immigrants (per generation)	N_R	1 – 10
Probability of recolonization (per generation)	P_R	0.05 – 0.1
Fecundity	F	5.5 ± 1.5

Table 2. Effects of demographic and genetic parameters (standardized effect size, given by the regression slope by its error) on time for adaptation (t_G) and on the mean-standardized selection coefficient (β_u) (see also Figs. S4 and S5 in Supplementary Online Material).

parameters	t_G	β_u
h^2	-18.5	12.9
cv	-11.7	10.1
v_U	-92.9	-11.2
O_0	22.6	-47.9
N_0	-79.2	-5.5
N_R	-49.2	12.9
P_R	-16.7	5.2
w^2	-1.3	38.2
F	-14.8	13.1
b	-3.6	58.3

For Review Only




Fig. 1. Distribution of time for adaptation (in generations) obtained from the IBM simulating dwarfing of *H. floresiensis* based on 10,000 random combinations of parameters defined in Table 1. Median time is equal to 346 generations (CI 95% ranging from 150 and 675 generations).




Fig. 2. Distribution of mean-standardized selection gradients generated in the IBM simulating dwarfing of *H. floresiensis* based on 10,000 random combinations of parameters defined in Table 1. Median selection strength is equal to -0.29 (CI 95% ranging from -0.167 and -0.562).